

Orbifold Hodge Numbers of Calabi-Yau Hypersurfaces

Mainak Poddar
University of Wisconsin-Madison
Email: poddar@math.wisc.edu

Abstract

We identify the twisted sectors of a compact simplicial toric variety. We do the same for a generic nondegenerate Calabi-Yau hypersurface of an n -dimensional simplicial Fano toric variety and then explicitly compute $h_{orb}^{1,1}$ and $h_{orb}^{n-2,1}$ for the hypersurface. We give applications to the orbifold string theory conjecture and orbifold mirror symmetry.

Contents

1	Introduction	2
2	Orbifolds	3
2.1	Orbifold Structure	3
2.2	Orbifold Cohomology	5
3	Facts from Toric Geometry	7
3.1	Fan, Orbits and Divisors	7
3.2	Projective Toric variety from polytope	9
3.3	Homogeneous coordinate ring	10
3.4	Fano toric Varieties	11
4	Twisted Sectors	12
4.1	Twisted Sectors in Simplicial Toric Variety	12
4.2	Twisted sectors of hypersurface of a Fano variety	14

5	Orbifold Hodge Numbers	15
5.1	$h_{orb}^{1,1}(V)$	15
5.2	$h^{n-2,1}(V)$	16
5.3	Cohomology of the twisted sectors of V	20
5.4	Main Results	22
5.5	An Example	23

1 Introduction

Recently, there is an emerging subject of mathematics: *Stringy geometry and topology of orbifolds*. The core of this includes the developing theories of orbifold cohomology, orbifold quantum cohomology, orbifold K-theory, motivic integration, McKay correspondence etc.

Physicists believe that orbifold string theory is equivalent to ordinary string theory of certain desingularisations. This belief motivated a body of conjectures, collectively referred to as the *Orbifold string theory conjecture*. The conjecture we are interested in is the *K-Orbifold string theory conjecture*. It states that there is a natural isomorphism between the *Orbifold K-theory* of a Gorenstein orbifold and the ordinary K-theory of its crepant resolution.

To construct a natural isomorphism as the conjecture demands, is a very hard problem. But many weaker versions of the conjecture that compare Euler numbers, Hodge numbers, etc. has been studied extensively in the literature in the case of orbifolds that are global-quotients. Batyrev[2], and Batyrev and Dais[4] proved, in particular, the equality of orbifold Hodge numbers and Hodge numbers of smooth crepant resolutions for Gorenstein global-quotient orbifolds. But there were no results for non global-quotient orbifolds.

In this paper, we show that the orbifold Hodge numbers of a generic Calabi-Yau hypersurface in a complex 4-dimensional simplicial Fano toric variety coincide with the Hodge numbers of its crepant resolution. Besides being the first non global-quotient example, this is also an important example in mirror symmetry. An immediate corollary of this is the mirror pairing of orbifold Hodge numbers for mirror families of Calabi-Yau 3-folds in simplicial Fano toric varieties. It should be remarked here that the orbifold structure on the Calabi-Yau hypersurface that we consider, is the one that naturally arises

from its algebraic structure.

Recently the author has observed that one can combine the characterisation of twisted sectors in this paper with the apparatus of string-theoretic Hodge numbers developed by Batyrev and Dais [4], and Batyrev [2] to greatly generalise the results of this paper in the toric world. There is even a hope that the problem may be completely solved, as far as Hodge numbers go, by using motivic integration [7] and some stack theory. But those results will be discussed elsewhere.

Now we briefly describe how this article is organised. In sections 2 and 3 we review some facts from orbifold cohomology and toric geometry respectively. In section 4 we find characterisations for the twisted sectors of simplicial toric varieties and nondegenerate Calabi-Yau hypersurfaces of simplicial Fano toric varieties. In section 5 we compute formulas for some orbifold Hodge numbers of these hypersurfaces, state some corollaries and then give an example.

Acknowledgements: This work was done under the supervision of my doctoral advisor, Prof Yongbin Ruan. I am indebted to him for his guidance. My special thanks are also due to Professors Igor Dolgachev, Lev Borisov and David Cox who were kind enough to listen to me and answer my questions. I thank Prof Cox for correcting an earlier draft of this article. I thank the faculty and friends at UW-Madison for their help and kindness.

2 Orbifolds

2.1 Orbifold Structure

Let U be a connected topological space, V a connected n -dimensional smooth manifold and G a finite group acting smoothly on V . An n -dimensional *uniformising system* of U is a triple (V, G, π) , where $\pi : V \rightarrow U$ is a continuous map inducing a homeomorphism between the quotient space V/G and U . Two uniformising systems $(V_i, G_i, \pi_i), i = 1, 2$, are *isomorphic* if there is a diffeomorphism $\phi : V_1 \rightarrow V_2$ and an isomorphism $\lambda : G_1 \rightarrow G_2$ such that ϕ is λ -equivariant, and $\pi_2 \circ \phi = \pi_1$.

Note: If (ϕ, λ) is an automorphism of (V, G, π) , then there is a $g \in G$

such that $\phi(x) = g.x$ and $\lambda(a) = g.a.g^{-1}$, for any $x \in V$ and $a \in G$.

Let $i : U' \rightarrow U$ be a connected open subset of U . An uniformising system (V', G', π') of U' is said to be induced from (V, G, π) if there is a monomorphism $\lambda : G' \rightarrow G$ and a λ -equivariant open embedding $\phi : V' \rightarrow V$ such that $i \circ \pi' = \pi \circ \phi$. The pair $(\phi, \lambda) : (V', G', \pi') \rightarrow (V, G, \pi)$ is called an *injection*. Two injections $(\phi_i, \lambda_i) : (V'_i, G'_i, \pi'_i) \rightarrow (V, G, \pi), i = 1, 2$, are *isomorphic* if there is an isomorphism (ψ, τ) from (V'_1, G'_1, π'_1) to (V'_2, G'_2, π'_2) , and an automorphism $(\bar{\psi}, \bar{\tau})$ of (V, G, π) such that $(\bar{\psi}, \bar{\tau}) \circ (\phi_1, \lambda_1) = (\phi_2, \lambda_2) \circ (\psi, \tau)$.

Fact: Let (V, G, π) be an uniformising system of U . For any connected open subset U' of U , (V, G, π) induces a unique isomorphism class of uniformising systems of U' .

Let U be a connected and locally connected topological space. For any point $p \in U$ we define the *germ* of uniformising systems at p in the following sense. Let (V_1, G_1, π_1) and (V_2, G_2, π_2) be uniformizing systems of neighbourhoods U_1 and U_2 of p . We say that (V_1, G_1, π_1) and (V_2, G_2, π_2) are *equivalent* at p if they induce isomorphic uniformising systems for a neighbourhood U_3 of p .

Definition: Let X be a Hausdorff, second countable topological space. An *n-dimensional orbifold structure* on X is given by the following data : for any point $p \in X$, there is a neighbourhood U_p and an n -dimensional uniformising system (V_p, G_p, π_p) of U_p such that for any point $q \in U_p$, (V_p, G_p, π_p) and (V_q, G_q, π_q) are equivalent at q .

Definition: The *germ of an orbifold structure* on X is an equivalence class under the following equivalence relation : two orbifold structures $\{(V_p, G_p, \pi_p) : p \in X\}$ and $\{(V'_p, G'_p, \pi'_p) : p \in X\}$ are equivalent if for any $p \in X$, (V_p, G_p, π_p) and (V'_p, G'_p, π'_p) are equivalent at p . With a given germ of orbifold structures on it, X is called an *orbifold*.

We call each U_p a uniformised neighbourhood of p , and (V_p, G_p, π_p) a chart at p . In fact we choose U_p to be small enough that G_p has the minimum possible order; that is, every element of G_p fixes the preimage of p in V_p . In what follows, this choice is assumed.

An open subset U of X is called an uniformised open set if it is uniformised by (V, G, π) such that for each $p \in U$, (V, G, π) defines the same germ as (V_p, G_p, π_p) at p . X is called a *global-quotient* orbifold if X itself is an uniformised open set. A point p is called *smooth* or *regular* if G_p is trivial; otherwise, it is called *singular*. The set of smooth points is denoted by X_{reg} , and the set of singular points is denoted by X_* .

An orbifold X is called **reduced** if G_p acts effectively on V_p . We can canonically associate a reduced orbifold to an orbifold by redefining the group actions. Furthermore if a group element acts nontrivially, we require that the fixed-point set is of at least (real) codimension two, so that the complement is locally connected. The toric varieties and hypersurfaces that we will deal with in this article are reduced orbifolds.

Remark: It is important to note here that even a reduced non-smooth orbifold can have a smooth underlying variety because of examples with complex reflections. For instance, consider $\mathbb{C}/\mathbb{Z}_2$ where $\mathbb{Z}_2$ acts by reflection about the origin. Then $\mathbb{C}/\mathbb{Z}_2$ is complex analytically isomorphic to $\mathbb{C}$ but it has a singularity at the origin in the orbifold sense. *Gorenstein* orbifolds do not present this problem as they do not admit such complex reflections.

2.2 Orbifold Cohomology

First we will describe the so-called twisted sectors. Consider the set of pairs:

$$\tilde{X}_* = \{(p, (g)_{G_p}) | p \in X_*, 1 \neq g \in G_p\},$$

where $(g)_{G_p}$ denotes the conjugacy class of g in G_p . Then Kawasaki showed (see [5]) that $\tilde{X}_*$ is naturally an orbifold with an orbifold structure given by

$$\{\pi_{p,g} : (V_p^g, C(g)/K_g) \rightarrow V_p^g/C(g) | p \in X_*, 1 \neq g \in G_p\},$$

where V_p^g is the fixed-point set of g in V_p , $C(g)$ is the centralizer of g in G_p , and K_g is the kernel of the action of $C(g)$ on V_p^g . There is a map $\pi : \tilde{X}_* \rightarrow X$ defined by $(p, (g)_{G_p}) \mapsto p$ such that $\pi(\tilde{X}_*) = X_*$. In a sense $\tilde{X}_*$ is a resolution of singularities of X .

Let $\tilde{X} = \tilde{X}_* \sqcup X$. We will describe the connected components of $\tilde{X}$. Recall that each point p has a local chart (V_p, G_p, π_p) which gives a local uniformized neighbourhood $U_p = \pi_p(V_p)$. If $q \in U_p$, up to conjugation, there is an injective homomorphism $G_q \rightarrow G_p$. For $g \in G_q$, the conjugacy class $(g)_{G_p}$ is well-defined. We define an equivalence relation $(g)_{G_q} \cong (g)_{G_p}$. Let T denote the set of equivalence classes. By an abuse of notation, we use (g) to denote the equivalence class to which $(g)_{G_q}$ belongs. $\tilde{X}$ is decomposed as a disjoint union of connected components

$$\tilde{X} = \bigsqcup_{(g) \in T} X_{(g)},$$

where

$$X_{(g)} = \{(p, (g')_{G_p}) | g' \in G_p, (g')_{G_p} \in (g)\}.$$

Definition: $X_{(g)}$ for $g \neq 1$ is called a **twisted sector**. We call $X_{(1)} = X$ the *untwisted* sector.

Let TX denote the *orbifold tangent bundle* of X . See Chen and Ruan[5] for a definition. An almost complex structure J on X is a smooth section of the orbifold bundle $End(TX)$ such that $J^2 = -Id$. Assume that X is an almost complex orbifold with an almost complex structure J . Then for any $p \in X_*$, J gives rise to an effective representation $\rho_p : G_p \rightarrow GL(n, \mathbb{C})$. For any $g \in G_p$ we write $\rho_p(g)$, upto conjugation, as a diagonal matrix $diag(e^{2\pi i \frac{m_{1,g}}{m_g}}, \dots, e^{2\pi i \frac{m_{n,g}}{m_g}})$, where m_g is the order of g in G_p , and $0 \leq m_{i,g} < m_g$. Define a function $\iota : \tilde{X} \rightarrow \mathbb{Q}$ by

$$\iota(p, (g)_{G_p}) = \sum_{i=1}^n \frac{m_{i,g}}{m_g}.$$

This function $\iota : \tilde{X} \rightarrow \mathbb{Q}$ is locally constant. Denote it by $\iota_{(g)}$. It has the following properties:

- (1) $\iota_{(g)}$ is integral iff $\rho_p(g) \in SL(n, \mathbb{C})$.
- (2) $\iota_{(g)} + \iota_{(g^{-1})} = rank(\rho_p(g) - Id) = n - dim_{\mathbb{C}} X_{(g)}$.

Definition: $\iota_{(g)}$ is called the *degree shifting number* of $X_{(g)}$.

Definition: An almost complex orbifold is called *Gorenstein* if property(1) holds for all (g) .

Remark: An almost complex, complex or kahler structure on X induces a corresponding similar structure on each $X_{(g)}$.

Definition: Let $\mathbb{F}$ be any field containing $\mathbb{Q}$ as a subfield. We define the *orbifold chomology groups* of X with coefficients in $\mathbb{F}$ by

$$H_{orb}^d(X; \mathbb{F}) = \oplus_{(g) \in T} H^{d-2\iota(g)}(X_{(g)}; \mathbb{F}).$$

Definition: Let X be a closed complex orbifold. We define, for $0 \leq p, q \leq \dim_{\mathbb{C}} X$, *orbifold Dolbeault cohomology groups*

$$H_{orb}^{p,q}(X; \mathbb{C}) = \oplus_{(g)} H^{p-\iota(g), q-\iota(g)}(X_{(g)}; \mathbb{C}).$$

Remark: When X is a closed Kahler orbifold (so is each $X_{(g)}$), these Dolbeault groups are related to the singular cohomology groups of X and $X_{(g)}$ as in the manifold case, and the Hodge decomposition theorem holds for these cohomology groups.

Definition: We define *orbifold Hodge numbers* by $h_{orb}^{p,q}(X) = \dim H_{orb}^{p,q}(X; \mathbb{C})$.

3 Facts from Toric Geometry

3.1 Fan, Orbits and Divisors

A complex n -dimensional toric variety is constructed from an n -dimensional lattice N and a fan Ξ in N . Ξ is a collection of strongly convex rational polyhedral cones σ in the real vector space $N_{\mathbb{R}} = N \otimes_{\mathbb{Z}} \mathbb{R}$, satisfying the conditons: every face of a cone in Ξ is also a cone in Ξ , and the intersection of two cones in Ξ is a face of each. We shall always assume that Ξ is finite.

Let $M = \text{Hom}(N, \mathbb{Z})$ denote the dual lattice, with dual pairing denoted by $\langle \cdot, \cdot \rangle$. If σ is a cone in N , the dual cone $\check{\sigma}$ is the set of vectors in $M_{\mathbb{R}}$ that are nonnegative on σ . This determines a commutative semigroup $S_{\sigma} = \check{\sigma} \cap M = \{u \in M : \langle u, v \rangle \geq 0 \text{ for all } v \in \sigma\}$. This semigroup is finitely generated. The corresponding group-algebra $\mathbb{C}[S_{\sigma}]$ which is the $\mathbb{C}$ -algebra with generators χ^m for each $m \in S_{\sigma}$ and relations $\chi^m \chi^{m'} = \chi^{m+m'}$, is a finitely generated commutative $\mathbb{C}$ -algebra. Such an algebra corresponds to an n -dimensional affine variety $U_{\sigma} := \text{spec} \mathbb{C}[S_{\sigma}]$.

If τ is a face of σ , then S_σ is contained in S_τ . So $\mathbb{C}[S_\sigma]$ is a subalgebra of $\mathbb{C}[S_\tau]$. This gives a map $U_\tau \rightarrow U_\sigma$. In fact, U_τ is a principal open subset of U_σ : if we choose $u \in S_\sigma$ so that $\tau = \sigma \cap u^\perp$, then $U_\tau = \{x \in U_\sigma : u(x) \neq 0\}$. With these identifications, these affine varieties for different cones fit together to form an algebraic variety X_Ξ . We will write X for X_Ξ when there is no confusion. X is a normal Cohen-Macaulay variety.

The properties of Ξ strongly affect the geometry of X . For example:

- (a) X is complete (i.e., compact) iff the *support* $|\Xi| := \cup_{\sigma \in \Xi} \sigma = N_\mathbb{R}$.
- (b) X is nonsingular iff for every cone in Ξ , its 1-dimensional generators (defined below) are part of a $\mathbb{Z}$ -basis of N . Such a fan is called *smooth*.
- (c) X is an orbifold iff the generators for every cone in Ξ are linearly independent over $\mathbb{R}$. We say X and Ξ are *simplicial*.

The affine variety corresponding to the trivial cone $\{0\}$ is the torus $T_N = N \otimes \mathbb{C} = \text{Spec} \mathbb{C}[M]$. It is an affine (Zariski) open subset of X_Ξ . The action of T_N on itself extends to an algebraic action of T_N on X_Ξ . This action has exactly one orbit corresponding to each cone $\tau \in \Xi$: $O_\tau = \text{Hom}(\tau^\perp \cap M, \mathbb{C}^*)$. We write $\tau < \sigma$ if τ is a face of σ and we write $\bar{O}_\sigma$ for the closure of O_σ in X . Then

- (a) $U_\sigma = \bigsqcup_{\tau < \sigma} O_\tau$; [Note $\tau = \sigma$ is included]
- (b) $\bar{O}_\tau = \bigsqcup_{\gamma > \tau} O_\gamma$;
- (c) $X_\Xi = \bigsqcup_{\tau \in \Xi} O_\tau$;

For each d , $\Xi(d)$ denotes the set of d dimensional cones of Ξ . We reserve the letter η to stand for elements of $\Xi(1)$. For each η , let v_η denote the unique generator of the semigroup $\eta \cap N$. Using these generators, a cone $\sigma \in \Xi$ can be written $\sigma = \{\sum_{\eta \subset \sigma} a_\eta v_\eta : a_\eta \geq 0\}$. The $v_\eta \in \sigma$ are the *generators* of σ . A standard abuse of notation is to identify η with v_η . If $r = |\Xi(1)|$ is the number of 1-dimensional cones, we sometimes write the v_η 's as $v_1, \dots, v_r$.

Each $\bar{O}_\eta$ is an irreducible T_N -invariant Weil divisor denoted D_η . Furthermore, $m \in M$ gives a character $\chi^m : T_N \rightarrow \mathbb{C}^*$, and regarding χ^m as a rational function on X , we have $\text{div}(\chi^m) = \sum_\eta \langle m, v_\eta \rangle D_\eta$. We will always assume that the 1-dimensional cones span $N_\mathbb{R}$. Then the Chow group $A_{n-1}(X)$ of Weil divisors modulo linear equivalence can be computed directly from the fan. By using the above, we get an exact sequence

$$0 \longrightarrow M \longrightarrow \mathbb{Z}^{\Xi(1)} \longrightarrow A_{n-1}(X) \longrightarrow 0$$

where $m \in M$ maps to $(\dots, \langle m, v_\eta \rangle, \dots) \in \mathbb{Z}^{\Xi(1)}$ and $(\dots, a_\eta, \dots) \in \mathbb{Z}^{\Xi(1)}$ maps to the divisor class of $\sum_\eta a_\eta D_\eta$. Thus $A_{n-1}(X)$ has rank $r - n$. A Weil divisor $D = \sum_\eta a_\eta D_\eta$ is Cartier iff for each $\sigma \in \Xi$, there is $m_\sigma \in M$ such that $\langle m_\sigma, v_\eta \rangle = -a_\eta$ whenever $\eta \subset \sigma$. Polytopes arise naturally when dealing with toric varieties. If X is complete and $D = \sum_\eta a_\eta D_\eta$ is Cartier, then

$$\begin{aligned} \Delta_D &= \{m \in M_{\mathbb{R}} : \langle m, v \rangle \geq \phi_D(v) \forall v \in N_{\mathbb{R}}\} \\ &= \{m \in M_{\mathbb{R}} : \langle m, v \rangle \geq -a_\eta \forall \eta\} \text{ is a polytope.} \end{aligned}$$

There is a T_N -equivariant map $H^0(X, \mathcal{O}(D)) \simeq \bigoplus_{m \in \Delta_D \cap M} \mathbb{C}\chi^m$. To see why this is true, think of sections of $\mathcal{O}(D)$ as rational functions f on X such that $\text{div}(f) + D \geq 0$. $\text{Div}(\chi^m) + D \geq 0$ is equivalent to $\langle m, v_\eta \rangle \geq -a_\eta \forall \eta$ and the isomorphism follows.

A Cartier divisor D is ample

$$\begin{aligned} &\iff \langle m_\sigma, v_\eta \rangle > -a_\eta \text{ whenever } \eta \text{ is not in } \sigma \text{ and } \sigma \text{ is } n\text{-dimensional.} \\ &\iff m_\sigma \neq m_\tau \text{ for } \sigma \neq \tau \text{ in } \Xi(n) \text{ and } \Delta_D \text{ is an } n\text{-dimensional polytope with} \\ &\text{vertices } \{m_\sigma : \sigma \in \Xi(n)\}. \end{aligned}$$

A polytope is called integral if its vertices are integral. Δ_D is integral if D is ample.

3.2 Projective Toric variety from polytope

Given any n -dimensional integral polytope Δ one can canonically associate a $\mathbb{C}$ -algebra S_Δ to it. See [[6], section 3.2.2] for details. Then the projective variety $\mathbb{P}_\Delta := \text{Proj}(S_\Delta)$ turns out to be a toric variety with a fan called the *normal fan* of Δ . We will describe the normal fan in the special case of Fano toric varieties later.

The T_N orbit closures of $\mathbb{P}_\Delta$ are in one-to-one correspondence with the nonempty faces F of Δ . There is a canonical surjection of polytope rings $\pi_F : S_\Delta \rightarrow S_F$ which induces a natural inclusion of toric varieties $\mathbb{P}_F \hookrightarrow \mathbb{P}_\Delta$. $\mathbb{P}_\Delta$ comes with a specific choice of ample divisor D_Δ such that $\Delta_{D_\Delta} = \Delta$.

Choose a basis for M . This corresponds to picking coordinates $t_1, \dots, t_n$ for the torus T_N . Then, if $m \in M$ is written $m = (a_1, \dots, a_n)$, we have $\chi^m = \prod_{i=1}^n t_i^{a_i}$, so we can write t^m instead of χ^m . For any $k \geq 0$, we have the space of Laurent polynomials $L(k\Delta) = \{f : f = \sum_{m \in k\Delta \cap M} \lambda_m t^m, \lambda_m \in \mathbb{C}\}$. Each $f \in L(k\Delta)$ gives the affine hypersurface $Z_f \subset T_N$ defined by $f = 0$. $L(k\Delta) \simeq H^0(\mathbb{P}_\Delta, \mathcal{O}_{\mathbb{P}_\Delta}(kD_\Delta))$. Under this isomorphism, f corresponds to an effective divisor $\bar{Z}_f \subset \mathbb{P}_\Delta$. $\bar{Z}_f$ is a compactification of Z_f and is exactly the hypersurface of $\mathbb{P}_\Delta$ corresponding to f . We will use the following notation:

- (a) $l(k\Delta) = |k\Delta \cap M| = \dim(L(k\Delta))$
- (b) $l^*(k\Delta) = |\{m \in k\Delta \cap M : m \text{ is not in any facet of } k\Delta \cap M\}|$

3.3 Homogeneous coordinate ring

We introduced coordinates $t_1, \dots, t_n$ on the torus of a toric variety. But it is useful to have global coordinates, similar to homogeneous coordinates on projective space. If X is given by a fan Ξ in $N_{\mathbb{R}}$, we introduce a variable x_η for each $\eta \in \Xi(1)$ and consider the polynomial ring $S = \mathbb{C}[x_\eta : \eta \in \Xi(1)]$. A monomial in S is written $x^D = \prod_\eta x_\eta^{a_\eta}$, where $D = \sum_\eta a_\eta D_\eta$ is an effective torus-invariant divisor on X . We say that x^D has degree $\deg(x^D) = [D] \in A_{n-1}(X)$.

Thus, S is graded by the Chow group $A_{n-1}(X)$. Given a divisor class $\alpha \in A_{n-1}(X)$, S_α denotes the graded piece of S of degree α . We often write the variables as $x_1, \dots, x_r$, where x_i corresponds to the cone generator v_i and $r = |\Xi(1)|$. Then $S = \mathbb{C}[x_1, \dots, x_r]$. The ring S , together with the grading defined above is called the *homogeneous coordinate ring* of X . We refer the reader to [[6], chapter 3.2] for a discussion on reconstructing the toric variety starting from the homogeneous coordinate ring.

If τ is any cone of Ξ then the orbit closure $\bar{O}_\tau$ is given by the ideal $(x_i : i \text{ such that } v_i \text{ is a generator of } \tau)$ of S . Also the graded pieces of S have nice cohomological interpretation. We noted that $L(\Delta) \simeq H^0(\mathbb{P}, \mathcal{O}_{\mathbb{P}}(D_\Delta))$. Now the map sending the Laurent monomial t^m to $\prod_\eta x_\eta^{\langle m, v_\eta \rangle + a_\eta}$ induces an isomorphism $H^0(X, \mathcal{O}_X(D)) \simeq S_\alpha$, where $\alpha = [D] \in A_{n-1}(X)$.

3.4 Fano toric Varieties

First we note that for an arbitrary toric variety X , the anticanonical divisor $-K_X = \sum_{\eta} D_{\eta}$. The following may be taken as definitions.

Definition: A toric variety X is *Gorenstein* iff $-K_X$ is Cartier. A complete toric variety X is *Fano* iff $-K_X$ is Cartier and ample.

The anticanonical divisor of a Fano toric variety X determines an integral polytope Δ with special properties :

- (a) All facets Γ of Δ are supported by an affine hyperplane of the form $\{m \in M_{\mathbb{R}} : \langle m, v_{\Gamma} \rangle = -1\}$ for some $v_{\Gamma} \in N$.
- (b) $\text{Int}(\Delta) \cap M = 0$.

Definition: Such a polytope is called *reflexive*.

The *polar polytope* Δ° of the reflexive polytope Δ is obtained by $\Delta^{\circ} = \{v \in N_{\mathbb{R}} : \langle m, v \rangle \geq -1 \text{ for all } m \in \Delta\} \subset N_{\mathbb{R}}$. The fan Ξ of X can be retrieved by coning over the proper faces of Δ° . This fan is called the *normal fan* of Δ and $X = \mathbb{P}_{\Delta}$. Δ° is also reflexive and $(\Delta^{\circ})^{\circ} = \Delta$. The Fano toric variety constructed from the normal fan of Δ° is denoted by $\mathbb{P}_{\Delta^{\circ}}$.

We shall use F and F° to denote a face of Δ and Δ° respectively. There exists an inclusion reversing duality between the faces of Δ and Δ° . For instance, the face of Δ which is dual to the face F° of Δ° is defined as $\widehat{F^{\circ}} := \{m \in \Delta : \langle n, m \rangle = -1 \forall n \in F^{\circ}\}$. If τ is the cone in the normal fan of Δ associated to the face F° , then the orbit closure $\bar{O}_{\tau} = \mathbb{P}_{\widehat{F^{\circ}}}$.

Generic anticanonical hypersurfaces V in $\mathbb{P}_{\Delta}$ and V° in $\mathbb{P}_{\Delta^{\circ}}$ constitute mirror families of **Calabi-Yau** varieties. These varieties are orbifolds if the corresponding ambient toric variety is simplicial. Observe that V can be identified with an element of $L(\Delta)$.

A ‘maximal projective subdivision’ of the fan of the toric variety gives a *maximal projective crepant partial* (MPCP) resolution [see [1] or [6]] of the hypersurface by taking proper transforms. Let $\widehat{V}$ and $\widehat{V}^{\circ}$ denote the MPCP resolutions of V and V° respectively. These are again Calabi-Yau. These are smooth if $n = 4$.

Remark: A gorenstein toric variety or a Calabi-Yau hypersurface of a

Fano toric variety has gorenstein canonical singularities. In particular, the degree shifting numbers are integers. Moreover, *the singularity locus has at least complex codimension two.*

4 Twisted Sectors

We claim that the twisted sectors of a toric variety or a Calabi-Yau hypersurface can be identified with suborbifolds. Note that in general a twisted sector could be a multiple cover of the corresponding singular locus even if the group actions are all abelian. The situation is simpler in toric case because there exist canonical local coordinates and uniformizing systems such that the group actions are in fact diagonal. They can be easily chased around to prove the claim.

4.1 Twisted Sectors in Simplicial Toric Variety

Let Ξ be any simplicial fan. Then the orbifold structure of the toric variety X_Ξ can be described as follows. Let σ be any n dimensional cone of Ξ . Let $v_1, \dots, v_n$ be the primitive 1 dim generators of σ . These are linearly independent in $N_{\mathbb{R}}$. Let N_σ be the sublattice of N generated by $v_1, \dots, v_n$. Let $G_\sigma := N/N_\sigma$ be the quotient group. G_σ is finite and abelian.

Let σ' be the cone σ regarded in N_σ . Let $\check{\sigma}'$ be the dual cone of σ' in M_σ , the dual lattice of N_σ . $U_{\sigma'} = \text{spec}(\mathbb{C}[\check{\sigma}' \cap M_\sigma])$. Note that σ' is a smooth cone in N_σ . So $U_{\sigma'} \cong \mathbb{C}^n$.

There is a canonical dual pairing $M_\sigma/M \times N/N_\sigma \rightarrow \mathbb{Q}/\mathbb{Z} \rightarrow \mathbb{C}^*$, the first map by the pairing $\langle, \rangle$ and the second by $q \mapsto \exp(2\pi i q)$. Now G_σ acts on $\mathbb{C}[M_\sigma]$, the group ring of M_σ , by $v(\chi^u) = \exp(2\pi i \langle u, v \rangle) \chi^u$, for $v \in N$ and $u \in M_\sigma$. Note that

$$(\mathbb{C}[M_\sigma])^{G_\sigma} = \mathbb{C}[M] \quad \text{.....(4.1.1)}$$

Thus G_σ acts on $U_{\sigma'}$. Let π_σ be the quotient map. Then $U_\sigma = U_{\sigma'}/G_\sigma$. So U_σ is uniformised by $(U_{\sigma'}, G_\sigma, \Pi_\sigma)$. For any $\tau < \sigma$, the orbifold structure on U_τ is same as the one induced from the uniformising system on U_σ . Then by the description of the toric gluing it is clear that $\{(U_{\sigma'}, G_\sigma, \Pi_\sigma) : \sigma \in \Xi(n)\}$ defines a reduced orbifold structure on X . We give a more explicit verification of this fact below.

Let $\mathcal{B}$ be the nonsingular matrix with generators $v_1, \dots, v_n$ of σ as rows. Then $\tilde{\sigma}'$ is generated in M_σ by the column vectors $v^1, \dots, v^n$ of the matrix $\mathcal{B}^{-1}$. So $\chi^{v^1}, \dots, \chi^{v^n}$ are the coordinates of $U_{\sigma'}$. For any $\kappa = (k_1, \dots, k_n) \in N$, the corresponding coset $[\kappa] \in G_\sigma$ acts on $U_{\sigma'}$ in these coordinates as a diagonal matrix: $\text{diag}(\exp^{2\pi i c_1}, \dots, \exp^{2\pi i c_n})$ where $c_i = \langle \kappa, v^i \rangle$. Such a matrix is uniquely represented by an n -tuple $a = (a_1, \dots, a_n)$ where $a_i \in [0, 1)$ and $c_i = a_i + b_i, b_i \in \mathbb{Z}$. In matrix notation, $\kappa \mathcal{B}^{-1} = a + b \iff \kappa = a\mathcal{B} + b\mathcal{B}$. We denote the integral vector $a\mathcal{B}$ in N by κ_a and the diagonal matrix corresponding to a by g_a . $\kappa_a \leftrightarrow g_a$ gives a one to one correspondence between the elements of G_σ and the integral vectors in N that are linear combinations of the generators of σ with coefficients in $[0, 1)$.

Now let us examine the orbifold chart induced by $(U_{\sigma'}, G_\sigma, \pi_\sigma)$ at any point $x \in U_\sigma$. By the orbit decomposition, there is unique $\tau \in \sigma$ such that $x \in O_\tau$. Wlog assume τ is generated by $v_1, \dots, v_j, j \leq n$. Then any preimage of x with respect to π_σ has coordinates $\chi^{v^i} = 0$ iff $i \leq j$. Let $z = (0, \dots, 0, z_{j+1}, \dots, z_n)$ be one such preimage. Let $G_\tau := \{g_a \in G_\sigma : a_i = 0 \text{ if } j+1 \leq i \leq n\}$. We can find a small neighbourhood $W \subset (\mathbb{C}^*)^{n-j}$ of $(z_{j+1}, \dots, z_n)$ such that the inclusions $\mathbb{C}^j \times W \hookrightarrow U_{\sigma'}$ and $G_\tau \hookrightarrow G_\sigma$ induces an injection of uniformising systems $(\mathbb{C}^j \times W, G_\tau, \pi) \hookrightarrow (U_{\sigma'}, G_\sigma, \pi_\sigma)$ on some small open neighbourhood U_x of x . So we have $G_x = G_\tau$ and an orbifold chart $(\mathbb{C}^j \times W, G_\tau, \pi)$. Note that G_τ can be constructed from the set $\{\kappa_a = \sum_{i=1}^j a_i V_i : \kappa_a \in N, a_i \in [0, 1)\}$ which is completely determined by τ and hence is independent of σ .

Now we determine the twisted sectors. Take any $x \in X$. x belongs to a unique O_τ . Wlog the generators of τ are $v_1, \dots, v_j$. Consider any n -dimensional $\sigma > \tau$. Wlog $v_1, \dots, v_n$ generates σ . Pick g_a in G_x such that $a_i \neq 0, \forall i \leq j$ i.e., κ_a lies in the interior of τ . We want to find the twisted sector $X_{(g_a)}$. Consider g_a as an element of G_σ . It is clear that g_a fixes $z \in U_{\sigma'}$ iff $z_1 = \dots = z_{j+s} = 0$, for some $s \geq 0$ i.e., $\pi_\sigma(z) \in O_\tau$ or $\pi_\sigma(z) \in O_\delta$ for some $\delta > \tau$. So $X_{(g_a)} \cap U_\sigma = \bar{O}_\tau \cap U_\sigma$. Since a twisted sector is connected, $X_{(g_a)} = \bar{O}_\tau$. If $g_a \in G_x$ is such that (wlog) only $a_1, \dots, a_k \neq 0, k < j$, then $g_a \in G_\delta$ where δ is the cone generated by $v_1, \dots, v_k$ and by the above argument $X_{(g_a)} = \bar{O}_\delta$. Thus we have proved the following theorem.

Theorem 1 *A twisted sector of any simplicial toric variety X_Ξ is isomorphic*

to a subvariety $\bar{O}_\tau$ of X_Ξ for some cone $\tau \in \Xi$. Moreover, there is a one-to-one correspondence between the set of twisted sectors of the type $\bar{O}_\tau$ and the set of integral vectors in the interior of τ which are linear combinations of the 1-dimensional generators of τ with coefficients in $(0, 1)$.

Note that the degree shifting number $\iota_{(g_a)} = \sum a_i$. Now if X_Ξ is Fano, i.e., Ξ is obtained by coning over the faces of a reflexive polytope Δ° , then the twisted sectors with $\iota = 1$ are in one to one correspondence with the integral interior points of faces of Δ° .

4.2 Twisted sectors of hypersurface of a Fano variety

We identify the twisted sectors of a generic nondegenerate anticanonical (Calabi-Yau) hypersurface V of a simplicial Fano toric variety $X = \mathbb{P}_\Delta$. **Nondegenerate** means that $V \cap O_\tau$ is either empty or a smooth subvariety of codimension one in O_τ , for each torus orbit O_τ in X . Then V turns out to be a suborbifold of X . Also nondegeneracy is a generic condition. For a different treatment on these, see [3]. We show that $V = \bar{Z}_f$, for a generic $f \in L(\Delta)$, is nondegenerate and a suborbifold of X .

Consider any n -dimensional cone σ with generators $v_1, \dots, v_n$. For notational simplicity set $\chi^{v^i} = z_i$. Then $z_1, \dots, z_n$ are the coordinates of $U_{\sigma'}$. Let Y be the preimage of $V \cap U_\sigma$ in $U_{\sigma'}$ with respect to π_σ . Then Y is defined by the equation $\sum_{m \in \Delta \cap M} \lambda_m \prod_{i=1}^n z_i^{\langle m, v_i \rangle + 1} = 0$. This is because, $t^m = \prod_{i=1}^n z_i^{q_i} \iff m = \sum q_i v^i = \mathcal{B}^{-1}q \iff q = m\mathcal{B} \iff q_i = \langle m, v_i \rangle$. The one is added to ensure that V is anticanonical. Note that by definition of Δ , $\langle m, v_i \rangle + 1 \geq 0$. If $\lambda_{m_\sigma} \neq 0$ then Y does not pass through the origin. It can be checked from this description using Bertini's theorem that for generic values of the coefficients λ_m , Y is a smooth submanifold of $U_{\sigma'}$ that intersects the coordinate planes $z_{i_1} = \dots = z_{i_j} = 0$ transversely.

Y is G_σ -stable by (4.1.1). When Y is smooth, all singularities of $V \cap U_\sigma$ are quotient singularities induced by action of G_σ on Y . Since there are only finitely many n -dimensional cones, V is nondegenerate and a suborbifold of X . $(Y, G_\sigma, \pi_\sigma)$ is a uniformizing system for $V \cap U_\sigma$. Let τ be the face of σ obtained by coning over the face F° of Δ° . Wlog let $v_1, \dots, v_j : j \geq 2$ be the generators of τ . (We remarked earlier in section 3.4 that the Gorenstein

condition rules out codimension one singularities.) We want to find a chart for any point $x \in V \cap O_\tau$. By our earlier remark that Y misses the origin of $U_{\sigma'}$, $V \cap \bar{O}_\sigma$ is empty. So we need only consider proper faces of σ . By a result of Fulton[[8],section 5.3], $V \cap \bar{O}_\tau$ consists of $l^*(\widehat{F^\circ}) + 1$ points if F° is a codimension 2 face of Δ° i.e., $j = n - 1 \iff \dim O_\tau = 1$. Since the only other points in $\bar{O}_\tau$ in this case are O_σ for n -dimensional cones $\sigma > \tau$, all the intersection points actually lie in O_τ . If codimension F° is bigger than 2, then again by Bertini $V \cap \bar{O}_\tau$ is irreducible.

Following the same notation as before, x has a small neighborhood $U_x \cap Y$ such that $((\mathbb{C}^j \times W) \cap Y, G_\tau, \pi)$ is a chart for V at x . $\mathbb{C}^j \times W$ is, as before, a suitable neighbourhood of some preimage z of x in $U_{\sigma'}$. The tangent space TY_z is a G_τ -stable subspace of $T\mathbb{C}_z^n$. Any $g_a \in G_\tau$ acts trivially on $TW_z = \text{span}\{\partial/\partial z_i, i = j + 1, \dots, n\}$. By transversality, we can choose basis $\{\xi_1, \dots, \xi_n\}$ of $T\mathbb{C}_z^n$ such that $\xi_i \in TY_z \forall i \leq n - 1$ and $\xi_n \in TW_z$. g_a acts trivially on ξ_n . This implies that the degree shifting number of $g_a|_{TY_z}$ is still $\sum_{i=1}^j a_i$.

From the description of the charts, it is clear that twisted sectors of V are isomorphic to $V \cap \bar{O}_\tau$ where $2 \leq \dim(\tau) \leq n - 1$. Recall that $\bar{O}_\tau = \mathbb{P}_{\widehat{F^\circ}}$ where $\widehat{F^\circ}$ is the face of Δ dual to F° . In particular we have the following theorem.

Theorem 2 *Let V be a generic nondegenerate anticanonical hypersurface of an n -dimensional simplicial Fano toric variety $\mathbb{P}_\Delta$. Then the twisted sectors of V are isomorphic to $V \cap \mathbb{P}_{\widehat{F^\circ}}$ for some face F° of Δ° such that $1 \leq \dim F^\circ \leq n - 2$. There is exactly one twisted sector of this type having $\iota = 1$, corresponding to each integral interior point of F° if $\dim F^\circ < n - 2$. If $\dim F^\circ = n - 2$, then there are exactly $l^*(\widehat{F^\circ}) + 1$ twisted sectors of this type having $\iota = 1$, corresponding to each integral interior point of F° .*

5 Orbifold Hodge Numbers

5.1 $h_{orb}^{1,1}(V)$

Let $V_{(g)}$ denote a twisted sector of the hypersurface V and $\iota_{(g)}$ its degree shifting number.

$h_{orb}^{1,1}(V) = h^{1,1}(V) + \sum_{\iota(g)=1} h^{0,0}(V_{(g)})$. Since $h^{0,0}(V_{(g)}) = 1$ for each twisted sector, by theorem (4.2) we obtain

$$\begin{aligned} & \sum_{\iota(g)=1} h^{0,0}(V_{(g)}) \\ &= \sum_{1 \leq \dim(F^\circ) \leq n-2} l^*(F^\circ) + \sum_{\dim(F^\circ)=n-2} l^*(F^\circ) l^*(\widehat{F^\circ}) \\ &= l(\Delta^\circ) - r - 1 - \sum_{\dim(F^\circ)=n-1} l^*(F^\circ) + \sum_{\dim(F^\circ)=n-2} l^*(F^\circ) l^*(\widehat{F^\circ}) \end{aligned}$$

To compute $h^{1,1}(V)$ we invoke the following Lefschetz hyperplane theorem [[3], Proposition 10.8] .

Lemma 1 *Let V be a nondegenerate ample hypersurface of an n -dimensional complete simplicial toric variety X . Then the natural map, induced by inclusion, $j^* : H^i(X) \rightarrow H^i(V)$ is an isomorphism for $i < n - 1$ and an injection for $i = n - 1$.*

In our case V is anticanonical, and since the anticanonical divisor of a Fano variety is ample, V is ample. Also it is well known[[8],section 5.1] that for any simplicial toric variety X , $H^2(X, \mathbb{R}) = H^{1,1}(X, \mathbb{R}) = A_{n-1}(X) \otimes \mathbb{R}$. So for $n \geq 4$, $h^{1,1}(V) = h^{1,1}(\mathbb{P}_\Delta) = \text{rank } A_{n-1}(\mathbb{P}_\Delta) = r - n$. Thus we have the following theorem.

Theorem 3 *For any generic nondegenerate anticanonical hypersurface V of an n -dimensional simplicial Fano toric variety $\mathbb{P}_\Delta$, $n \geq 4$,*

$$h_{orb}^{1,1}(V) = l(\Delta^\circ) - n - 1 - \sum_{\dim(F^\circ)=n-1} l^*(F^\circ) + \sum_{\dim(F^\circ)=n-2} l^*(F^\circ) l^*(\widehat{F^\circ}).$$

5.2 $h^{n-2,1}(V)$

Next we compute $h^{n-2,1}(V)$ for $n \geq 4$. For this we have to use the homogeneous coordinate ring S of $X = \mathbb{P}_\Delta$. Let $v_1, \dots, v_r$ be the one dimensional cones of the normal fan of Δ . Let $x_1, \dots, x_r$ be the corresponding homogeneous coordinates. Let $\beta_0 = [-K_X] = [\sum_{i=1}^r D_i] \in A_{n-1}(X)$. Then $S_{\beta_0} \simeq L(\Delta)$. And the divisor $f \in L(\Delta)$ corresponding to V can be written in the homogeneous coordinates as $f = \sum_{m \in M \cap L(\Delta)} \lambda_m \prod_{i=1}^r x_i^{\langle m, v_i \rangle + 1}$. For notational simplicity, we will denote $\prod_{i=1}^r x_i^{\langle m, v_i \rangle + 1}$ by $\mathbf{x}^m$ for any $m \in M$. Define the Jacobian ideal of f to be $J(f) = \langle \partial f / \partial x_1, \dots, \partial f / \partial x_r \rangle$.

First we quote the following theorem [[3], theorem 10.13] .

Lemma 2 *Let X be an d -dimensional complete simplicial toric variety and $V \subset X$ be a quasi-smooth (i.e., suborbifold) ample hypersurface defined by $f \in S_\beta$. Then for $k \neq (d/2) + 1$, there exists a canonical isomorphism*

$$(S/J(f))_{k\beta-\beta_0} \simeq PH^{d-k,k-1}(V).$$

Remark: The primitive cohomology $PH^{d-1}(V) := H^{d-1}(V)/(\text{im} H^{d-1}(X))$. This coincides with the usual cohomology if d is even.

For our application of this lemma: $d = n, k = 2$ and $\beta = \beta_0$. So we need to compute the rank of $(S/J(f))_{\beta_0}$. Also, $H^{n-2,1}(X) = 0$ if $n \geq 4$, so that $H^{n-2,1}(V) = PH^{n-2,1}(V)$.

Lemma 3 *$x_i \partial f / \partial x_i \in (J(f))_{\beta_0}, i = 1, \dots, r$, and the space of complex linear relations among these has dimension $r - (n + 1)$.*

Proof

$$\begin{aligned} x_i \partial f / \partial x_i &= \sum_m \lambda_m (\langle m, v_i \rangle + 1) \mathbf{x}^m. \\ \sum_i c_i x_i \partial f / \partial x_i &= \sum_m \lambda_m (\langle m, \sum_i c_i v_i \rangle + \sum_i c_i) \mathbf{x}^m \end{aligned}$$

For a generic f we can assume that $\lambda_m \neq 0$ for each $m \in \Delta \cap M$. Hence $\sum_i c_i x_i \partial f / \partial x_i \equiv 0 \iff \langle m, \sum_i c_i v_i \rangle + \sum_i c_i = 0 \forall m \in \Delta \cap M$. In particular, taking $m = 0$ we get $\sum_i c_i = 0$. Therefore $\langle m, \sum_i c_i v_i \rangle = 0 \forall m \in \Delta \cap M$ and since Δ is n -dimensional we have $\sum_i c_i v_i = 0$. Thus $\sum_i c_i x_i \partial f / \partial x_i \equiv 0 \iff \sum_i c_i v_i = 0$ and $\sum_i c_i = 0$.

Now let $\tilde{v}_i = (v_i, 1) \in \mathbb{R}^{n+1} \subset \mathbb{C}^{n+1} = \mathbb{R}^{n+1} \otimes \mathbb{C}$.

Since the v_i 's are vertices of Δ° , the $\tilde{v}_i$'s are generators of the $(n + 1)$ -dimensional cone $\{q \in \mathbb{R}^{n+1} : qt \in \Delta^\circ \text{ for some } t \in \mathbb{R}_{>0}\}$. So the $\tilde{v}_i$'s span $\mathbb{R}^{n+1}$ over $\mathbb{R}$, and hence they span $\mathbb{R}^{n+1} \otimes \mathbb{C}$ over $\mathbb{C}$.

Note that $(\sum c_i v_i, \sum c_i) = \sum c_i \tilde{v}_i$.

Hence the lemma follows.

Wlog assume that the $x_k \partial f / \partial x_k, k = 1, \dots, n + 1$, are linearly independent. In other words $\tilde{v}_k, k = 1, \dots, n + 1$ are linearly independent. We want to find monomials $\prod_{j \neq i} x_j^{p_j}$ that have same degree in S as x_i . So we want $m^* \in M$ such that $\langle m^*, v_j \rangle = p_j \geq 0$ if $j \neq i$ and $\langle m^*, v_i \rangle = -1$. Such m^* is given by

interior lattice points of F_i , the $(n-1)$ -dimensional face of Δ that is dual to the 0-dimensional face $\{v_i\}$ of Δ° .

Then for each $m^* \in (\text{Int} F_i \cap M)$, $\prod_{j \neq i} x_j^{\langle m^*, v_j \rangle} \partial f / \partial x_i$ belongs to $(J(f))_{\beta_0}$. Together with the $x_i \partial f / \partial x_i$, these generate $(J(f))_{\beta_0}$ as we vary over all i .

$$\begin{aligned}
& \prod_{j \neq i} x_j^{\langle m^*, v_j \rangle} \partial f / \partial x_i \\
&= \prod_{j=1}^r x_j^{\langle m^*, v_j \rangle} x_i \partial f / \partial x_i \\
&= \sum_{m \in \Delta \cap M} \lambda_m (\langle m, v_i \rangle + 1) \mathbf{x}^{m+m^*} \\
&= \sum_{m' \in \Delta \cap M \cap (\Delta + m^*)} \lambda_{m'-m^*} (\langle m' - m^*, v_i \rangle + 1) \mathbf{x}^{m'} \\
&= \sum_{m \in \Delta \cap M} \lambda_{m-m^*} (\langle m - m^*, v_i \rangle + 1) I(m - m^* \in \Delta) \mathbf{x}^m
\end{aligned}$$

Let $\text{Int} F_i \cap M = \{m_{i,i_s} : 1 \leq s \leq t_i; t_i \geq 0\}$.

Then $(J(f))_{\beta_0} = \text{span}\{x_k \partial f / \partial x_k, \prod_{j=1}^r x_j^{\langle m_{i_s}, v_j \rangle} x_i \partial f / \partial x_i : 1 \leq k \leq n+1, 1 \leq i_s \leq t_i, i = 1, \dots, r\}$. We want to find the dimension of this complex vector space. So we study the space of linear relations :

$$\begin{aligned}
& \sum_k c_k x_k \partial f / \partial x_k + \sum_{i,i_s} d_{i,i_s} \prod_{j=1}^r x_j^{\langle m_{i_s}, v_j \rangle} x_i \partial f / \partial x_i \equiv 0 \\
& \iff \\
& \sum_m \left\{ \sum_k c_k \lambda_m (\langle v_k, m \rangle + 1) + \sum_{i,i_s} d_{i,i_s} \lambda_{m-m_{i,i_s}} I(m-m_{i,i_s} \in \Delta) (\langle m-m_{i,i_s}, v_i \rangle + 1) \right\} \mathbf{x}^m \equiv 0 \\
& \iff \\
& \sum_k c_k \lambda_m (\langle v_k, m \rangle + 1) + \sum_{i,i_s} d_{i,i_s} \lambda_{m-m_{i,i_s}} I(m-m_{i,i_s} \in \Delta) \langle m, v_i \rangle \equiv 0 \text{ for each } m \in \Delta \cap M, \\
& \hspace{25em} [\text{note: } \langle m_{i,i_s}, v_i \rangle = -1]
\end{aligned}$$

This is a system of $l(\Delta)$ number of linear equations in $\gamma = n+1 + \sum_{i=1}^r l^*(F_i)$ variables namely c_k, d_{i,i_s} . Note that $l(\Delta) \geq \gamma$. We shall find a nonsingular subsystem of rank γ .

To do so pick n linearly independent vertices $m_1, \dots, m_n$ of Δ and let $m_{n+1} = 0$, the origin. Then from the above system we pick the equations corresponding to $m = m_1, \dots, m_{n+1}$ and $m = m_{i,i_s} : i = 1, \dots, r; 0 \leq i_s \leq t_i$.

Denote this $\gamma \times \gamma$ system by (**). It can be written as :

$$\begin{bmatrix} \mathbf{P} & \mathbf{A} \\ \mathbf{B} & \mathbf{Q} \end{bmatrix} \begin{pmatrix} c \\ d \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

where

$$\mathbf{P} = \begin{bmatrix} \lambda_{m_1}(\langle m_1, v_1 \rangle + 1) & \dots & \lambda_{m_1}(\langle m_1, v_{n+1} \rangle + 1) \\ \dots & \dots & \dots \\ \lambda_{m_n}(\langle m_n, v_1 \rangle + 1) & \dots & \lambda_{m_n}(\langle m_n, v_{n+1} \rangle + 1) \\ \lambda_0 & \dots & \lambda_0 \end{bmatrix}$$

$$\mathbf{Q} = \begin{bmatrix} \lambda_{m_{1,1}-m_{1,1}} I(\cdot)(\langle m_{1,1} - m_{1,1}, v_1 \rangle + 1) & \dots & \lambda_{m_{1,1}-m_{r,t_r}} I(\cdot)(\langle m_{1,1} - m_{r,t_r}, v_r \rangle + 1) \\ \dots & \dots & \dots \\ \lambda_{m_{r,t_r}-m_{1,1}} I(\cdot)(\langle m_{r,t_r} - m_{1,1}, v_1 \rangle + 1) & \dots & \lambda_{m_{r,t_r}-m_{r,t_r}} I(\cdot)(\langle m_{r,t_r} - m_{r,t_r}, v_r \rangle + 1) \end{bmatrix}$$

Observe that all the diagonal entries of $\mathbf{Q}$ are λ_0 , and none of its off-diagonal entries has λ_0 . Also any entry of $\mathbf{A}$ is of the form $\lambda_{m_k-m_{i,i_s}} I(\cdot)$ and hence does not involve λ_0 . Similarly an entry of $\mathbf{B}$ is of the form $\lambda_{m_{i,i_s}}(\langle m_{i,i_s}, v_k \rangle + 1)$ and so does not have λ_0 .

Consider the determinant of the coefficient matrix $\begin{bmatrix} \mathbf{P} & \mathbf{A} \\ \mathbf{B} & \mathbf{Q} \end{bmatrix}$ as a polynomial in the λ 's. Then the term of this determinant having the highest power of λ_0 is $(\lambda_0)^{\sum l^*(F_i)} \det \mathbf{P}$. We will show below that $\det \mathbf{P} = \text{nonzero constant}$ times $\lambda_{m_1} \dots \lambda_{m_n} \lambda_0$. Thus the determinant of the coefficient matrix of the system (**) is a nontrivial polynomial in the λ 's and is therefore nonzero for generic choice of the λ 's. Hence $(J(f))_{\beta_0}$ has rank γ as a complex vector space, for a generic $f \in L(\Delta)$. Since $S_{\beta_0} \simeq L(\Delta)$, so $(S/J(f))_{\beta_0}$ has rank $l(\Delta) - \gamma$, for a generic f .

Lemma 4 *The $(n+1) \times (n+1)$ matrix $\mathbf{P} = ((P_{i,j} = \lambda_{m_i}(\langle m_i, v_j \rangle + 1)))$ is nonsingular for generic choice of λ 's.*

Proof

Let $\mathbf{E}$ be the $(n+1) \times (n+1)$ matrix $((E_{i,j} = (\langle m_i, v_j \rangle + 1)))$. Then $\det \mathbf{P} = \lambda_{m_1} \dots \lambda_{m_{n+1}} \det \mathbf{E}$.

We claim that $\mathbf{E}$ is nonsingular. Otherwise there exists a nontrivial vector $(c_1, \dots, c_{n+1})$ such that $\sum_{k=1}^{n+1} c_k(\langle m_i, v_k \rangle + 1) = 0$ for all $i = 1, \dots, n+1$. In particular, for $i = n+1$ we get $\sum_{k=1}^{n+1} c_k = 0$. This implies $\sum_{k=1}^{n+1} c_k \langle m_i, v_k \rangle = 0$ for all $i = 1, \dots, n$. Since $m_1, \dots, m_n$ are linearly independent, this would imply that $\sum_{k=1}^{n+1} c_k \langle m, v_k \rangle = 0$ for all $m \in \Delta$. Therefore $\sum_{k=1}^{n+1} c_k v_k = 0$. This combined with $\sum c_k = 0$ implies that $\sum_{k=1}^{n+1} c_k \tilde{v}_k = 0$ which contradicts the linear independence of $\tilde{v}_1, \dots, \tilde{v}_{n+1}$. Thus the lemma holds.

So we have the following theorem.

Theorem 4 *For any generic nondegenerate anticanonical hypersurface V of an n -dimensional simplicial Fano toric variety $\mathbb{P}_\Delta$, and $n \geq 4$,*

$$h^{n-2,1}(V) = l(\Delta) - n - 1 - \sum_{\dim(F)=n-1} l^*(F).$$

Note : F denotes a facet of the polytope Δ in the statement of the above theorem.

5.3 Cohomology of the twisted sectors of V

We now want to compute $h^{n-3,0}(V_{(g)})$ for any twisted sector $V_{(g)} \cong V \cap \mathbb{P}_{\widehat{F^\circ}}$. This is obviously zero if $\dim(F^\circ) > 1$, since $\dim(\mathbb{P}_{\widehat{F^\circ}}) = n - 1 - \dim(F^\circ)$. So we will only consider the case $\dim(F^\circ) = 1$. Let τ be the 2-dimensional cone obtained by coning over F° . As noted earlier $\bar{O}_\tau = \mathbb{P}_{\widehat{F^\circ}}$. The restriction of V to $\bar{O}_\tau$ gives a quasismooth ample hypersurface of $\bar{O}_\tau$, which we shall identify with $V_{(g)}$. So we are again in a situation where we can invoke lemma 2.

For this we need to understand the homogeneous coordinate ring S' of $\bar{O}_\tau$. According to Fulton[[8],section 3.1], a fan for $\bar{O}_\tau$ can be constructed from the fan Ξ of X as follows.

Let N_τ be the sublattice of N generated by the primitive one dimensional generators of τ . Let $N(\tau) = N/N_\tau$. The dual lattice of $N(\tau)$ is given by $M(\tau) = \tau^\perp \cap M$. The *star* of a cone τ can be defined abstractly as the set of cones σ in Ξ that contain τ as a face. Such cones σ are determined by their images in $N(\tau)$ i.e. by $\bar{\sigma} = (\sigma + (N_\tau)_\mathbb{R}) / (N_\tau)_\mathbb{R} \subset N_\mathbb{R} / (N_\tau)_\mathbb{R} = N(\tau)_\mathbb{R}$. These cones $\{\bar{\sigma} : \tau < \sigma\}$ form a fan $\text{Star}(\tau)$ in $N(\tau)$. $\bar{O}_\tau$ is the toric variety corresponding to this fan.

Wlog let v_1, v_2 be the generators of τ . The corresponding Weil divisors in X are D_1 and D_2 , and $\bar{O}_\tau$ is the intersection $|G_\tau|D_1D_2$ considered as an element of the Chow ring of X .

Again wlog assume that $v_j, j = 3, \dots, l$ are the one dimensional cones of Ξ such that $\{v_1, v_2, v_j\}$ generate a three dimensional cone of Ξ . In other words, $\bar{v}_j, j = 3, \dots, l$ are the one dimensional cones of $\text{Star}(\tau)$. Let $\tilde{D}_j := D_1D_2D_j$ for $j = 3, \dots, r$. Note that $\tilde{D}_j = 0$ if $j > l$. The divisor of $\bar{O}_\tau$ corresponding to $\bar{v}_j$ is $\tilde{D}_j$ for $j = 3, \dots, l$. So the homogeneous coordinate ring S' is generated by variables y_j corresponding to $\tilde{D}_j$ for $j = 3, \dots, l$. Denote by α_0 the anti-canonical class $\sum_{j=3}^l \tilde{D}_j$ in S' .

On the other hand the divisor V restricts to $-K_X D_1 D_2 = (D_1 + \dots + D_r) D_1 D_2 = \sum_{j=3}^l \tilde{D}_j + (D_1 + D_2) D_1 D_2$. To see what $(D_1 + D_2) D_1 D_2$ is in terms of the $\tilde{D}_j$ s, we can pick a point $m \in \widehat{F^\circ} \cap M$ and let $b_i := \langle m, v_i \rangle, 1 \leq i \leq r$. Then $\sum_{i=1}^r b_i D_i$ is linearly equivalent to zero. Note that $b_1 = b_2 = -1$. Hence we have $D_1 + D_2 = \sum_{i=3}^r b_i D_i$. So, $(D_1 + D_2) D_1 D_2 = \sum_{i=3}^l b_i \tilde{D}_i$. Let α be the class in S' representing the effective ample divisor $-K_X D_1 D_2$. Let f' be the associated homogeneous polynomial in the y_j s. Now we can apply lemma 2 to the $(n-2)$ -dimensional variety $\bar{O}_\tau$ and the ample hypersurface $V_{(g)}$. Choose $k = 1$ in the lemma to get

$$(S'/J(f'))_{\alpha-\alpha_0} \simeq PH^{n-3,0}(V_{(g)}) = H^{n-3,0}(V_{(g)})$$

since $H^{n-3,0}(\bar{O}_\tau) = 0$.

Now $\alpha - \alpha_0 = [\alpha - \sum_{j=3}^l \tilde{D}_j]$. A typical generator $\partial f' / \partial y_i \in J(f')$ has degree $[\alpha - \tilde{D}_i]$. There are no nonconstant regular functions on the projective variety $\bar{O}_\tau$. So any nontrivial effective divisor, and in particular $\tilde{D}_i, \sum_{j=3, \dots, l; j \neq i} \tilde{D}_j$ and $\sum_{j=3, \dots, l} \tilde{D}_j$ are not linearly equivalent to zero. This implies that $(J(f'))_{\alpha-\alpha_0} = 0$. Hence we obtain that

$$(S')_{\alpha-\alpha_0} \simeq H^{n-3,0}(V_{(g)})$$

Now $\alpha - \alpha_0 = [\sum_{j=3}^l b_j \tilde{D}_j]$. We want to identify the effective divisors in this class. So we want $m_* \in M(\tau)$ such that $\sum_{j=3}^l (b_j + \langle m_*, \bar{v}_j \rangle) \tilde{D}_j$ is effective. This is if and only if $(b_j + \langle m_*, \bar{v}_j \rangle) \geq 0$ for all $j = 3, \dots, l$

$$\iff (\langle m + m_*, v_j \rangle) \geq 0 \text{ for } j = 3, \dots, l$$

$$\iff m + m_* \in \widehat{\text{Int} F^\circ} \cap M.$$

To justify the last step note that $\langle m + m_*, v_i \rangle = -1$ for $i = 1, 2$ so that $m + m_* \in \widehat{F^\circ} \cap M$. If $m + m_*$ is not in the interior of $\widehat{F^\circ}$ then it belongs to some face $\widehat{E^\circ}$ where $F^\circ \subset E^\circ$ as faces of Δ° . The cone δ associated to E° contains τ as a face and hence contains some v_k for $3 \leq k \leq l$. Then $\langle m + m_*, v_k \rangle = -1$ which is a contradiction.

Since m is fixed, the required effective divisors are in one-to-one correspondence with the interior lattice points of $\widehat{F^\circ}$. Hence $h^{n-3,0}(V_{(g)}) = l^*(\widehat{F^\circ})$. Since there are $l^*(F^\circ)$ twisted sectors isomorphic to $V \cap \mathbb{P}_{\widehat{F^\circ}}$ we have the following.

$$\begin{aligned} h_{orb}^{n-2,1}(V) &= h^{n-2,1}(V) + \sum_{\iota(g)=1} h^{n-3,0}(V_{(g)}) \\ &= l(\Delta) - n - 1 - \sum_{\dim(F)=n-1} l^*(F) + \sum_{\dim(F^\circ)=1} l^*(F^\circ) l^*(\widehat{F^\circ}) \\ &= l(\Delta) - n - 1 - \sum_{\dim(F)=n-1} l^*(F) + \sum_{\dim(F)=n-2} l^*(F) l^*(\widehat{F}). \end{aligned}$$

For the last step we used the one-to-one correspondence between faces of Δ and Δ° .

5.4 Main Results

Combining the above formula with theorem 4 we have the following theorem.

Theorem 5 *For any generic nondegenerate anticanonical hypersurface V of an n -dimensional simplicial Fano toric variety $\mathbb{P}_\Delta$, $n \geq 4$,*

$$h_{orb}^{n-2,1}(V) = l(\Delta) - n - 1 - \sum_{\dim(F)=n-1} l^*(F) + \sum_{\dim(F)=n-2} l^*(F) l^*(\widehat{F}).$$

Corollary 1 *If $\widehat{V}$ is an MPCP desingularisation of any generic nondegenerate anticanonical hypersurface V of an n -dimensional simplicial Fano toric variety $\mathbb{P}_\Delta$, $n \geq 4$, then $h_{orb}^{p,1}(V) = h^{p,1}(\widehat{V})$ for $p = 1$ and $p = n - 2$*

Proof The formulas for $h^{p,1}(\widehat{V})$ for $p = 1, n - 2$ computed in [1] by Batyrev match the orbifold Hodge numbers for V obtained in theorem 3 and theorem 5.

Corollary 2 *In the case $n = 4$, $h_{orb}^{p,q}(V) = h^{p,q}(\widehat{V})$ for any p and q .*

Proof We need only consider $p, q \leq 3$. Also $h_{orb}^{p,0} \equiv h^{p,0}$ by definition since ι is nonnegative. So, by Serre duality, it is enough to consider just the cases $p = 1, q = 1$ and $p = 2, q = 1$. These are addressed by corollary 1.

We should remark here that in this case $\widehat{V}$ is actually smooth.

Corollary 3 *If $\mathbb{P}_{\Delta^\circ}$ is also simplicial, and V° is a generic nondegenerate anticanonical hypersurface of $\mathbb{P}_{\Delta^\circ}$, then $h_{orb}^{1,1}(V) = h_{orb}^{n-2,1}(V^\circ)$ and vice versa.*

Proof Follows from interchanging the roles of Δ and Δ° in the formulas.

Remark In particular, for the $n = 4$ case, we have $h_{orb}^{p,q}(V) = h_{orb}^{3-p,q}(V^\circ)$. This is an example of ‘mirror symmetry’ of orbifold hodge numbers.

5.5 An Example

Consider the complex 4-dimensional weighted projective space $X = \mathbb{P}(1,1,2,2,2)$. It is a simplicial Fano toric variety. Its fan Ξ has the following 1-dimensional cones in $N \cong \mathbb{Z}^4$: $v_1 = (-1, -2, -2, -2)$, $v_2 = (1, 0, 0, 0)$, $v_3 = (0, 1, 0, 0)$, $v_4 = (0, 0, 1, 0)$, $v_5 = (0, 0, 0, 1)$. Ξ has five 4-dimensional cones, obtained by dropping one of the v_i ’s at a time and taking the cone generated by the remaining four.

Let D_i denote the torus-invariant divisor given by the orbit closure $\overline{O_{v_i}}$. It is easy to check that in $A_3(X)$, $[D_2] = [D_1]$ and $[D_i] = 2[D_1]$ for $i \geq 3$. Construct the homogeneous coordinate ring of X by introducing variables x_i corresponding to v_i . Then $\deg(x_1) = \deg(x_2) = 1$ ($= [D_1]$) and $\deg(x_i) = 2$ for $i \geq 3$.

This leads to the more familiar description of $\mathbb{P}(1,1,2,2,2)$ as $(\mathbb{C}^5 - \{0\})/\mathbb{C}^*$. The action of any $\alpha \in \mathbb{C}^*$ on $\mathbb{C}^5 - \{0\}$ is as follows :
 $\alpha \cdot [x_1 : x_2 : x_3 : x_4 : x_5] = [\alpha x_1 : \alpha x_2 : \alpha^2 x_3 : \alpha^2 x_4 : \alpha^2 x_5]$.

In this description, D_i corresponds to the hyperplane $\{x_i = 0\}$ and the 4-dimensional cones of Ξ correspond to the open sets $\{x_i \neq 0\}$. It is also easily seen that the singular locus of X is precisely the surface $\{x_1 = x_2 = 0\}$. In fact, this represents the only twisted sector of X . The v_i ’s are the vertices of a reflexive polytope Δ° . The faces of Δ° have only one interior lattice point : $(0, -1, -1, -1) = 1/2 (v_1 + v_2)$. This lattice point corresponds to the twisted

sector and the local isotropy group is $\mathbb{Z}_2$.

The dual reflexive polytope Δ in $M_{\mathbb{R}}$ has the following vertices:
 $w_1 = (-1, -1, -1, -1)$, $w_2 = (7, -1, -1, -1)$, $w_3 = (-1, 3, -1, -1)$,
 $w_4 = (-1, -1, 3, -1)$, $w_5 = (-1, -1, -1, 3)$.

Δ is the polytope corresponding to the anticanonical divisor $-K_X = \sum_{i=1}^5 D_i$ of X , and $X = \mathbb{P}_{\Delta}$. If V is a generic nondegenerate Calabi-Yau (anticanonical) hypersurface of X , then V has just one twisted sector namely $V \cap \{x_1 = x_2 = 0\}$. One can directly compute the genus of this curve by using the Riemann-Hurwitz formula. It turns out to be 3.

The following is a list of the Hodge numbers of V : $h^{1,0} = h^{2,0} = 0$, $h^{3,0} = 1$, $h^{1,1} = 1$, $h^{2,1} = 83$, $h_{orb}^{1,1} = 2$, $h_{orb}^{2,1} = 86$.

The dual Fano variety $\mathbb{P}_{\Delta^\circ}$ is also simplicial. This is easily checked since its fan is obtained by coning over the faces of Δ . In fact, $\mathbb{P}_{\Delta^\circ} = \mathbb{P}_{\Delta}/\mathbb{Z}_4^3$. This is also shown easily. First, observe that $w_1 = -w_2 - 2w_3 - 2w_4 - 2w_5$. Secondly, if $\bar{M}$ is the sublattice of M generated by w_2, w_3, w_4, w_5 , then $M/\bar{M} = \mathbb{Z}_4^3$.

The mirror Calabi-Yau family is also easy to write down. Let z_i denote the homogeneous coordinate of $\mathbb{P}_{\Delta^\circ}$ corresponding to w_i . Then a generic Calabi-Yau hypersurface V° of $\mathbb{P}_{\Delta^\circ}$ has equation:

$$\lambda_1 z_1^8 + \lambda_2 z_2^8 + \lambda_3 z_3^4 + \lambda_4 z_4^4 + \lambda_5 z_5^4 + \lambda_6 z_1^4 z_2^4 + \lambda_0 z_1 z_2 z_3 z_4 z_5 = 0.$$

Note that $h_{orb}^{1,1}(V^\circ) = 86$ and $h_{orb}^{2,1}(V^\circ) = 2$.

Bibliography

- [1] V.Batyrev, *Dual polyhedra and Mirror symmetry for Calabi-Yau hypersurfaces in toric varieties*, AG/9310003
- [2] V.Batyrev, *Non-Archimedean Integrals and Stringy Euler numbers of log-terminal pairs*, JEMS 1, 1999.
- [3] V.Batyrev and D.Cox, *On the Hodge structure of projective hypersurfaces in toric varieties*, AG/9306011

- [4] V.Batyrev and D.Dais, *Strong McKay correspondence, string-theoretic Hodge numbers and mirror symmetry*, AG/9410001
- [5] W.Chen and Y.Ruan, *A New cohomology theory for Orbifold*, AG/0004129
- [6] D.Cox and S.Katz, *Mirror Symmetry and Algebraic Geometry*, Mathematical Surveys and Monographs, Vol 68, AMS.
- [7] J.Denef and F.Loester, *Germes of arcs on singular algebraic varieties and motivic integration*, Invent. Math. **135** (1999).
- [8] W.Fulton, *Introduction to toric varieties*, Princeton University Press, 1993.
- [9] M.Reid, *La correspondance de McKay*, AG/9911165
- [10] Y.Ruan, *Stringy geometry and Topology of Orbifolds*, AG/0011149